



Research

Cite this article: Hare JF, Campbell KL, Senkiw RW. 2014 Catch the wave: prairie dogs assess neighbours' awareness using contagious displays. *Proc. R. Soc. B* **281**: 20132153. <http://dx.doi.org/10.1098/rspb.2013.2153>

Received: 17 August 2013

Accepted: 3 December 2013

Subject Areas:

behaviour, cognition, ecology

Keywords:

contagious communication, group-size effect, public information, theory of mind, foraging–vigilance trade-off, prairie dogs

Author for correspondence:

James F. Hare

e-mail: james.hare@ad.umanitoba.ca

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rspb.2013.2153> or via <http://rspb.royalsocietypublishing.org>.

Catch the wave: prairie dogs assess neighbours' awareness using contagious displays

James F. Hare, Kevin L. Campbell and Robert W. Senkiw

Department of Biological Sciences, University of Manitoba, Winnipeg, Manitoba, Canada R3T 2N2

The jump yip display of black tailed prairie dogs (*Cynomys ludovicianus*) is contagious, spreading through a prairie dog town as 'the wave' through a stadium. Because contagious communication in primates serves to assess conspecific social awareness, we investigated whether instigators of jump yip bouts adjusted their behaviour relative to the response of conspecifics recruited to display bouts. Increased responsiveness of neighbouring town members resulted in bout initiators devoting a significantly greater proportion of time to active foraging. Contagious jump yips thus function to assess neighbours' alertness, soliciting social information to assess effective conspecific group size in real time and reveal active probing of conspecific awareness consistent with theory of mind in these group living rodents.

1. Introduction

Members of group living species rely upon collective vigilance to detect predators [1], occasionally synchronizing vigilance so as to enhance the detection of potential threats [2,3]. As the number of individuals residing within a group increases, each individual can devote less time to vigilance and more time to other activities, for example foraging [1,4,5], owing, in part, to the cumulative effect of each individual's finite probability of detecting presumptive threats [6–8]. Where such group size effects are dependent upon collective detection, however, individual group members must assess the size of the group in which they reside [9], and perhaps more importantly, the vigilance of fellow group members [10,11] to tailor their behaviour to the situation at hand.

Black tailed prairie dogs (*Cynomys ludovicianus*) are group living, semi fossorial rodents that are subject to intense predation by terrestrial and avian predators [12,13]. Classic group size effects, with diminishing individual allotment of time to vigilance and increasing time devoted to foraging with increasing group size, have been documented observationally [14] and experimentally [15] among black tailed prairie dogs, though the mechanism via which group size is assessed is unknown. Just as synchrony in the activity of group members promotes social cohesion [16,17] through social facilitation among individuals [18], fine tuning one's foraging vigilance trade off relative to the vigilance of other group members would prove adaptive [11].

We investigated contagious, multimodal 'jump yip' displays of black tailed prairie dogs as a potential means of adjusting the instigating individual's foraging vigilance trade off relative to the responsiveness, and hence vigilance, of neighbouring group members. In instigating a jump yip bout, an individual raises its anterior torso above the ground, achieving at least an erect posture, though sometimes propelling itself from the ground. This postural change is coupled with the extension of the signaller's forelimbs and the emission of a vocalization that can be described phonetically as 'wee oo' [19] as the anterior torso is raised and then lowered, resulting in a multimodal display lasting around a second ([20]; see the electronic supplementary material, movie S1). Unlike other prairie dog vocalizations or displays, the jump yip is typically contagious, spreading from the initiator through neighbouring colony members as 'the wave' through a stadium ([20]; see the electronic supplementary material, movie S2). To date, the function of the jump yip has proved elusive.

Initial accounts were purely descriptive [21,22], though subsequent researchers have suggested that the jump yip serves as a territorial call [12], an ‘all clear’ [13,23] or ‘end of danger’ signal [12], a means of promoting social bonding within the group (‘contact group cohesion’; [19]), an indicator that the signaller is less likely to show escape behaviour in the face of threat, promoting temporarily heightened vigilance among conspecifics [20], or as Owings & Coss [24] suggested for various animal signals, a means of managing conspecifics. Here, such management would presumably promote reliance on conspecifics for predator detection, thereby allowing the signaller to devote time to activities other than watchfulness.

The acquisition of information from conspecifics proves adaptive in contexts including foraging [25], habitat choice [26], mate choice [27], parenting [28], tool use [29], and both anti brood parasite [30] and anti predator responses [31]. Although social learning and public information use are often considered tacitly different concepts [32–34], the effective utilization of socially available information requires that individuals assess the veracity of the information they receive [35–37], and sample available information on an ongoing basis to ensure that information is up to date [38]. Such assessment is of particular importance where ignoring information proves costly [39], as would be the case where animals are faced with the risk of predation [8].

We hypothesized that the instigation of a jump yip bout serves to probe neighbouring individuals for feedback regarding their current vigilance state. If that were the case, we predicted that subsequent time devoted to personal vigilance by bout instigators would be directly proportional to the latency of the first conspecific to respond to the instigator’s jump yip (greater delay in response indicating reduced alertness of neighbours), and inversely proportional to the number of conspecifics joining a jump yip bout (a smaller number indicating reduced collective vigilance) as well as the overall duration of the bout (reduced length of contagion indicative of lesser collective vigilance). Given the well documented trade off between foraging and vigilance in black tailed prairie dogs [13–15], we also predicted that the allocation of time to foraging by bout instigators would be inversely correlated with the latency of the first conspecific to respond in kind with a jump yip and positively correlated with the number of individuals recruiting to, and the overall duration of jump yip bouts.

2. Material and methods

To ascertain whether instigator vigilance and foraging behaviour were affected by subsequent jump yip bout properties (latency to first response, number of individuals recruited and bout duration), we video recorded a total of 173 jump yip bouts during November 2003 and from May through September 2004 within 16 distinct prairie dog towns spread across six populations, including 14 naturally occurring towns in South and North Dakota, USA, and two introduced towns in Winnipeg, Manitoba, Canada (see R.L. Senkiw M.Sc. thesis for further details; downloadable at: <http://mspace.lib.umanitoba.ca/bitstream/1993/2842/1/MSc%20Thesis.pdf>). To ensure our results were unfounded, we eliminated bouts interrupted by the appearance of predators, prairie dog vocalizations other than those constituting part of the jump yip bout, humans or vehicles, and those where the behaviour of the bout instigator could not be distinguished because it was out of frame or out of focus in the video recording, or where any ambiguity existed regarding the identity of the bout initiator. Thus, our final dataset included data from

27 independent bout initiators from 14 distinct towns (one to three bouts per town) among the six populations, for which we quantified the proportion of time the bout instigator engaged in vigilance (with its head above the horizontal plane) and the proportion of time spent foraging (head below the horizontal plane, grazing or chewing) in the 1 min subsequent to the second syllable of the bout instigator’s ‘wee oo’ call. Postures in which the head was below the horizontal plane but where neither active grazing nor chewing occurred were not scored as vigilance or foraging. Limiting estimation of vigilance and foraging to only 1 min maximized the likelihood that the bout instigator’s behaviour was attributable solely to properties of the current jump yip bout. Bouts were considered independent of each other if at least 4 s elapsed without an individual manifesting a jump yip, based on the 5 s criterion Smith *et al.* [20] employed to delineate unique bouts, and on an obvious discontinuity in the distribution of individuals recruiting to bouts after 4 s in our larger 173 bout sample of videotaped jump yip bouts. For each bout, we recorded the latency (s) for the first individual to respond with a jump yip to the bout instigator, the number of jump yip responses within each bout and the overall duration of each bout (s). We applied linear regressions to test for relationships between those three independent variables, and the proportions of time the bout instigator devoted to either vigilance or foraging in the post bout period, considering those statistically significant where $p < 0.05$.

3. Results

The proportion of time devoted to vigilance by bout initiators was unaffected by the latency for the initial respondent to recruit to a contagious jump yip bout ($F_{1,25} = 0.03$, $p = 0.87$; figure 1). Further, although time allocated to vigilance tended to decrease with both an increasing number of respondents and increasing bout duration, those declines fell short of statistical significance ($F_{1,25} = 2.86$, $p = 0.10$ and $F_{1,25} = 2.89$, $p = 0.10$, respectively; figure 1). Similarly, although there was a trend toward decreasing time devoted to foraging with increasing latency of conspecifics to respond to the initial propagation of a jump yip, that change was not statistically significant ($F_{1,25} = 1.31$, $p = 0.26$; figure 1). Statistically significant relationships were detected for the proportionate allocation of time to active foraging, relative to both the number of respondents recruiting to a given jump yip bout ($F_{1,25} = 5.87$, $p = 0.02$) and the overall duration of contagious display bouts ($F_{1,25} = 4.50$, $p = 0.04$). Bout instigators increased time allocated to foraging with both an increasing number of individuals recruiting to a jump yip bout and increasing bout duration (figure 1). That said, bout duration was positively correlated with the number of respondents ($r = 0.82$), though neither bout duration nor the number of respondents recruiting to a jump yip bout was well correlated with latency to response ($r = 0.32$ and $r = 0.09$, respectively). Significant differences or pronounced trends for bout initiators to alter their allocation of time to foraging and vigilance in accord with our predictions exist for five of the six relationships examined, which in itself, should occur rarely by chance alone (binomial $p = 0.11$).

4. Discussion

Instigators of jump yip bouts increased their proportionate allocation of time to foraging as the responsiveness of conspecifics increased. The absence of any relationship between




Figure 1. Proportionate allocation of black-tailed prairie dog jump–yip bout instigator time to foraging and vigilance relative to the latency (s) of the first respondent to contribute to a contagious bout of calling, the number of respondents recruiting to the bout and the overall duration (s) of the jump–yip bout evoked (* $p < 0.05$).

bout instigator time allocation to vigilance and latency of initial conspecific response is not attributable to bout instigators consistently manifesting maximal vigilance, in that the proportion of time devoted to vigilance varied considerably across the range of response latencies observed. Moreover, bout instigators are undoubtedly constrained to devote time to vigilance so as to assess bout response characteristics prior to adjusting their foraging–vigilance trade off accordingly. The effect of conspecific responsiveness on the proportion of time devoted to foraging is not attributable to variation in group size proper, in that treating aboveground group size as a covariate did not alter our findings. Jump–yip displays thus function to promote the accrual of information regarding collective vigilance within the group, clarifying earlier speculation regarding the function of these displays.

While conspecific responsiveness to the instigation of a jump–yip bout significantly affected the time bout initiators allocated to foraging, only an inconsequential change was detected in instigator personal vigilance with variation in conspecific response. How accurately our simple postural assay of the head being held above the horizontal plane reflects an individual's state of vigilance remains an open question, as does consideration of any potential cost associated with the use of this socially acquired information [40].

Our results are consistent with the assertion by Owings & Coss [24] that tonic communication functions to manage conspecifics, in this case providing up to date, context specific information on the vigilance of conspecific group members. Among group living species, accurate decision making is facilitated via the use of social information [41,42], particularly where certain group members are less well informed than others [43]. Conspecific neighbours constitute a particularly important source of information where animals are subject to predation risk [44], and reliance on that information may, in and of itself, prompt social contagion [45]. In mimicking the evasive behaviour of neighbours, group living insects avoid unseen

predators (the 'Trafalgar effect'; [46]). Similarly, schooling fish avoid previously unencountered predators by mimicking their neighbours' behaviour [47].

Based on both previous empirical findings and an individual based model, Beauchamp *et al.* [11] argued that selection would favour monitoring and copying the vigilance of neighbours, resulting in collective waves of vigilance that would facilitate collective detection of predators. Given the coevolutionary nature of predator–prey relationships, however, predators might be expected to cue in on, and coordinate their attacks relative to lulls in such waves. Our empirical findings for black-tailed prairie dogs reveal collective waves of coordinated behaviour, presumably optimizing foraging efficiency of individuals reciprocally sharing information regarding collective vigilance. Participants within a bout thus likely benefit via reciprocal altruism [48] among resident town members, with any risk of enhanced detection by presumptive predators owing to the production of this conspicuous display being shared among signallers. Such broadly subscribed patterns of display behaviour may also serve as a potent pronouncement of vigilance [49], thereby reducing predation risk among town members in general.

The wave-like spread of jump–yip displays through towns of black-tailed prairie dogs is consistent with literature implicating the use of public information in social evolution [33,34] and contagious displays in particular, in the evaluation of conspecific social awareness [50]. Yawning in humans and other primates [51,52] and laughter among humans [53] provide familiar examples of contagious displays. Platek *et al.* [51], among others, have reported that such behaviours are associated with self-processing and empathy in humans. Indeed, emotional contagion among humans has been considered at least an important precursor to more advanced 'Theory of mind' abilities [54], wherein contagion represents the first step toward respondent awareness of the emotional state of the instigator by invoking that same state in the

respondent [55]. Emotional contagion in humans, however, transcends superficial motor mimicry and represents a means of sharing affect of considerable evolutionary antiquity [56]. Although the existence of a relationship between empathy and social contagion has long been recognized for non human primates [57], it has more recently been reported for dogs [58]. Applied to our findings, it is evident that in responding to the emergent display properties of neighbours recruited to a jump yip bout, black tailed prairie dogs manifest at least a rudimentary awareness of the state of conspecific group members. As Barrett *et al.* [59] aptly point out, communication and mind are intimately intertwined, in that the 'mind' is a form of social participation or process, rather than an entity unto itself, ultimately serving to rationalize the organism's social environment. In this light, it is not surprising that these highly social animals have evolved coordinated social behaviour and commensurate cognitive abilities [60] promoting their success in the face of intense predation pressure. Further study is required to elucidate additional nuances of the mechanism, along with the ultimate implications of this newly discovered, adaptive social contagion in prairie dogs.

All research adhered to the guidelines of the Canadian Council on Animal Care as detailed under Protocol 03 034 approved by the University of Manitoba's Fort Garry Campus Protocol Management and Review Committee, and under US National Park Service Scientific Research Permit THRO 2003 SCI 0011.

Acknowledgements. We thank Drs Daniela Campobello, Maurizio Sarà, Debbie Kelly and two anonymous reviewers, along with Associate Editor Dr Nadia Aubin Horth for insightful comments on an earlier draft of this manuscript. We also thank Dr Susan Cosens for input during the course of R.S.'s thesis work, and Dr Bob Wrigley (Assini boine Park Zoo), Ken Cudmore (Fort Whyte Centre), Penny Knuckles and Michael Oehler (Theodore Roosevelt National Park), Terry Lincoln (Dakota Zoo), Dan Miller (Bramble Park Zoo) and Dan Foster (Wind Cave National Park) for facilitating access to the prairie dogs.

Data accessibility. Bout response characteristic and bout initiator behavioural data file: Dryad doi:10.5061/dryad.g76t8.

Funding statement. Funding for this research was provided by a University of Manitoba Graduate Fellowship and a Sigma Xi Grant in Aid of Research to R.S., and by the Natural Sciences and Engineering Research Council of Canada in the form of a Postgraduate Scholarship to R.S. and Discovery grant (no. 154271) funding to J.H.

References

- Elgar M. 1989 Predator vigilance and group size in mammals and birds: a critical review of the empirical evidence. *Biol. Rev.* **64**, 13–33. (doi:10.1111/j.1469-185X.1989.tb00636.x)
- Pays O, Renaud PC, Loisel P, Petit M, Gerard J, Jarman PJ. 2007 Prey synchronize their vigilant behavior with other group members. *Proc. R. Soc. B* **274**, 1287–1291. (doi:10.1098/rspb.2006.0204)
- Townsend SW, Zöttl M, Manser M. 2011 All clear? Meerkats attend to contextual information in close calls to coordinate vigilance. *Behav. Ecol. Sociobiol.* **65**, 1927–1934. (doi:10.1007/s00265-011-1202-6)
- Quenette P Y. 1990 Functions of vigilance behaviour in mammals: a review. *Acta Oecol.* **11**, 801–818.
- Roberts G. 1996 Why individual vigilance declines as group size increases. *Anim. Behav.* **51**, 1077–1086. (doi:10.1006/anbe.1996.0109)
- Pulliam HR. 1973 On the advantages of flocking. *J. Theor. Biol.* **38**, 419–422. (doi:10.1016/0022-5193(73)90184-7)
- Lima SL. 1995 Back to the basics of anti predatory vigilance: the group size effect. *Anim. Behav.* **49**, 11–20. (doi:10.1016/0003-3472(95)80149-9)
- Lima SL, Dill LM. 1990 Behavioral decisions made under the risk of predation: review and prospectus. *Can. J. Zool.* **68**, 619–640. (doi:10.1139/z90-092)
- Elgar MA, Burren PJ, Posen M. 1984 Vigilance and perception of flock size in foraging house sparrows (*Passer domesticus* L.). *Behaviour* **90**, 215–223. (doi:10.1163/156853984X00146)
- Lima SL, Zollner PA. 1996 Anti predatory vigilance and the limits to collective detection: visual and spatial separation between foragers. *Behav. Ecol. Sociobiol.* **38**, 355–363. (doi:10.1007/s00265005-0252)
- Beauchamp G, Alexander P, Jovani R. 2012 Consistent waves of collective vigilance in groups using public information about predation risk. *Behav. Ecol.* **23**, 368–374. (doi:10.1093/beheco/arr194)
- King JA. 1955 Social behavior, social organization, and population dynamics in a black tailed prairie dog town in the Black Hills of South Dakota. *Contrib. Lab. Vert. Biol.* **67**, 1–123.
- Hoogland JL. 1995 *The black tailed prairie dog: social life of a burrowing mammal*. Chicago, IL: University of Chicago Press.
- Hoogland JL. 1979 The effect of colony size on individual alertness of prairie dogs: Sciuridae (*Cynomys* spp.). *Anim. Behav.* **27**, 394–407. (doi:10.1016/0003-3472(79)90174-X)
- Kildaw SD. 1995 The effect of group size manipulations on the foraging behavior of black tailed prairie dogs. *Behav. Ecol.* **6**, 353–358. (doi:10.1093/beheco/6.4.353)
- Focardi S, Pecchiol E. 2005 Social cohesion and foraging decrease with group size in fallow deer (*Dama dama*). *Behav. Ecol. Sociobiol.* **59**, 84–91. (doi:10.1007/s00265-005-0012-0)
- Gautrais J, Michelena P, Sibbald A, Bon R, Deneubourg J L. 2007 Allelomimetic synchronization in Merino sheep. *Anim. Behav.* **74**, 1443–1454. (doi:10.1016/j.anbehav.2007.02.020)
- Clayton DA. 1978 Socially facilitated behavior. *Q. Rev. Biol.* **53**, 373–392. (doi:10.1086/410789)
- Waring GH. 1970 Sound communications of black tailed, white tailed, and Gunnison's prairie dogs. *Am. Midl. Nat.* **83**, 167–185. (doi:10.2307/2424014)
- Smith WJ, Smith SL, DeVilla JG, Oppenheimer EC. 1976 The jump yip display of the black tailed prairie dog, *Cynomys ludovicianus*. *Anim. Behav.* **24**, 609–621. (doi:10.1016/S0003-3472(76)80075-9)
- Seton ET. 1926 The prairie dogs (*Cynomys ludovicianus*) at Washington Zoo. *J. Mamm.* **7**, 229–230.
- Anthony A. 1955 Behavior patterns in a laboratory colony of prairie dogs, *Cynomys ludovicianus*. *J. Mamm.* **36**, 69–78. (doi:10.2307/1375724)
- Smith RE. 1958 *Natural history of the prairie dog in Kansas*. Lawrence, KS: University of Kansas Press.
- Owings DH, Coss RG. 2007 *Social and antipredator systems: intertwining links in multiple time frames*. In *Rodent societies: ecological and evolutionary perspectives* (eds J Wolff, PW Sherman), pp. 305–316. Chicago, IL: University of Chicago Press.
- Giraldeau L A, Caraco T. 2000 *Social foraging theory*. Princeton, NJ: Princeton University Press.
- Danchin E, Boulinier T, Massot M. 1998 Conspecific reproductive success and breeding habitat selection: implications for the study of coloniality. *Ecology* **79**, 2415–2428. (doi:10.2307/176832)
- Nordell RR, Valone T. 1998 Mate choice copying as public information. *Ecol. Lett.* **1**, 74–76. (doi:10.1046/j.1461-0248.1998.00025.x)
- Forsman JT, Seppänen JT, Nykänen IL. 2012 Observed heterospecific clutch size can affect offspring investment decisions. *Biol. Lett.* **8**, 341–343. (doi:10.1098/rsbl.2011.0970)
- Whiten A, Horner V, de Waal FBM. 2005 Conformity to cultural norms of tool use in chimpanzees. *Nature* **437**, 737–740. (doi:10.1038/nature04047)
- Campobello D, Sealy SG. 2011 Use of social over personal information enhances nest defense against avian brood parasitism. *Behav. Ecol.* **22**, 422–428. (doi:10.1093/beheco/arq225)
- Cornell HN, Marzluff JM, Pecoraro S. 2012 Social learning spreads knowledge about dangerous humans among American crows. *Proc. R. Soc. B* **279**, 499–508. (doi:10.1098/rspb.2011.0957)
- Bonnie KE, Earley RL. 2007 Expanding the scope for social information use. *Anim. Behav.* **74**, 171–181. (doi:10.1016/j.anbehav.2006.12.009)

33. Danchin E, Giraldeau L A, Valone TJ, Wagner RH. 2004 Public information: from nosy neighbors to cultural evolution. *Science* **305**, 487–491. (doi:10.1126/science.1098254)
34. Dall SRX, Giraldeau L A, Olsson O, McNamara JM, Stephens DW. 2005 Information and its use by animals in evolutionary ecology. *Trends Ecol. Evol.* **20**, 187–193. (doi:10.1016/j.tree.2005.01.010)
35. Cheney DL, Seyfarth RM. 1988 Assessment of meaning and the detection of unreliable signals by vervet monkeys. *Anim. Behav.* **36**, 477–486. (doi:10.1016/S0003-3472(88)80018-6)
36. Hare JF, Atkins BA. 2001 The squirrel that cried wolf: reliability detection by juvenile Richardson's ground squirrels. *Behav. Ecol. Sociobiol.* **51**, 108–112. (doi:10.1007/s002650100414)
37. Searcy WA, Nowicki A. 2005 *The evolution of animal communication: reliability and deception in signaling systems*. Princeton, NJ: Princeton University Press.
38. van Bergen Y, Coolen I, Laland KN. 2004 Nine spined sticklebacks exploit the most reliable source when public and private information conflict. *Proc. R. Soc. Lond. B* **271**, 957–962. (doi:10.1098/rspb.2004.2684)
39. Wray MK, Klein BA, Seeley TD. 2012 Honey bees use social information in waggle dances more fully when foraging errors are more costly. *Behav. Ecol.* **23**, 125–131. (doi:10.1093/beheco/arr165)
40. Giraldeau L A, Valone TJ, Templeton JJ. 2002 Potential disadvantages of using socially acquired information. *Phil. Trans. R. Soc. Lond. B* **357**, 1559–1566. (doi:10.1098/rstb.2002.1065)
41. Seppänen J T, Forsman JT, Mönkkönen M, Thomson RL. 2007 Social information use is a process across time, space, and ecology, reaching heterospecifics. *Ecology* **88**, 1622–1633. (doi:10.1890/06-1757.1)
42. Farrell S. 2011 Social influence benefits the wisdom of individuals in the crowd. *Proc. Natl Acad. Sci. USA* **108**, E625. (doi:10.1073/pnas.1109947108)
43. King AJ, Cowlshaw G. 2007 When to use social information: the advantage of large group size in individual decision making. *Biol. Lett.* **3**, 137–139. (doi:10.1098/rsbl.2007.0017)
44. Fernández Juricic E, Beauchamp G, Treminio R, Hoover M. 2011 Making heads turn: association between head movements during vigilance and perceived predation risk in brown headed cowbird flocks. *Anim. Behav.* **82**, 573–577. (doi:10.1016/j.anbehav.2011.06.014)
45. Sirost E. 2006 Social information, antipredatory vigilance and flight in bird flocks. *Anim. Behav.* **72**, 373–382. (doi:10.1016/j.anbehav.2005.10.028)
46. Treherne JE, Foster WA. 1981 Group transmission of predator avoidance behaviour in a marine insect: the Trafalgar affect. *Anim. Behav.* **29**, 911–917. (doi:10.1016/S0003-3472(81)80028-0)
47. Brown C, Laland KN. 2003 Social learning in fishes: a review. *Fish. Fish.* **4**, 280–288. (doi:10.1046/j.1467-2979.2003.00122.x)
48. Trivers RL. 1971 The evolution of reciprocal altruism. *Q. Rev. Biol.* **46**, 35–57. (doi:10.1086/406755)
49. Hasson O. 1991 Pursuit deterrent signals: communication between prey and predators. *Trends Ecol. Evol.* **6**, 325–329. (doi:10.1016/0169-5347(91)90040-5)
50. Hecht EE, Patterson R, Barbey AK. 2012 What can other animals tell us about human social cognition? An evolutionary perspective on reflective and reflexive processing. *Front. Hum. Neurosci.* **6**, 224. (doi:10.3389/fnhum.2012.00224)
51. Platek SM, Critton SR, Myers TE, Gallup Jr GG. 2003 Contagious yawning: the role of self awareness and mental state attribution. *Cogn. Brain Res.* **17**, 223–237. (doi:10.1016/S0926-6410(03)00109-5)
52. Anderson JR, Myowa Yamakoshi M, Matsuzawa T. 2004 Contagious yawning in chimpanzees. *Proc. R. Soc. Lond. B* **271**, S468–S470. (doi:10.1098/rsbl.2004.0224)
53. Provine RR. 1996 Laughter. *Am. Sci.* **84**, 38–45.
54. Gallese V, Keysers C, Rizzolatti G. 2004 A unifying view of the basis of social cognition. *Trends Cogn. Sci.* **8**, 396–403. (doi:10.1016/j.tics.2004.07.002)
55. Preston SD, de Waal FBM. 2002 Empathy: its ultimate and proximate bases. *Behav. Brain Sci.* **25**, 1–71. (doi:10.1017/S0140525X02000018)
56. Tamietto M, Castelli L, Vighetti S, Perozzo P, Geminiani G, Weiskrantz L, de Gelder B. 2009 Unseen facial and bodily expressions trigger fast emotional reactions. *Proc. Natl Acad. Sci. USA* **106**, 17 661–17 666. (doi:10.1073/pnas.0908994106)
57. Brothers L. 1990 The neural basis of primate social communication. *Motiv. Emotion* **14**, 81–91. (doi:10.1007/BF00991637)
58. Silva K, de Sousa L. 2011 'Canis empathicus': a proposal on dogs' capacity to empathize with humans. *Biol. Lett.* **7**, 489–492. (doi:10.1098/rsbl.2011.0083)
59. Barrett L, Henzi SP, Lusseau D. 2012 Taking sociality seriously: the structure of multi dimensional social networks as a source of information for individuals. *Phil. Trans. R. Soc. B* **367**, 2108–2118. (doi:10.1098/rstb.2012.0113)
60. Slobodchikoff CN, Perla BS, Verdolin JL. 2009 *Prairie dogs: communication and community in an animal society*. Cambridge, MA: Harvard University Press.