

THE UNIVERSITY OF MANITOBA

DATE: April 14, 2003

FINAL EXAMINATION

PAPER NO: 131

PAGE 1 of 8

DEPARTMENT & COURSE NO: 136.130

TIME: 2 HOURS

EXAMINATION: Vector Geometry &
Linear Algebra

EXAMINERS: Various

Values

[10] 1. Let $A = \begin{pmatrix} 1 & 0 & 2 \\ 2 & 1 & 1 \\ 1 & 1 & 2 \end{pmatrix}$, $B = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix}$, $C = \begin{pmatrix} 1 & 3 \\ -2 & -1 \\ 1 & 2 \end{pmatrix}$.

Either evaluate each of the following expressions or give a reason why it is not defined.

- (a) $2B - 3C^T$
- (b) CAB
- (c) $A(C+B)$
- (d) $\text{tr}(BC) - \text{tr}(CB)$

[9] 2. Find all the values of b for which the following system of equations:

$$\begin{aligned}x + 2y &= b \\ 2x + (3 + b^2)y &= 3b + 1\end{aligned}$$

- (a) has no solutions;
- (b) has infinitely many solutions;
- (c) has a unique solution.

[10] 3. Express the matrix $M = \begin{pmatrix} 6 & -2 \\ -5 & 2 \end{pmatrix}$ as a product of elementary matrices.

[12] 4. Let A be the coefficient matrix of the following system of equations.

$$\begin{aligned}2x + 6y - 2z &= 1 \\ y - z &= 2 \\ x + 4y - z &= 3\end{aligned}$$

- (a) Write the system in the form of a matrix equation.
- (b) Find the cofactor C_{21} for the matrix A .
- (c) Find A^{-1} .
- (d) Solve the system using part 4(d) above.

[12] 5. Let $M = \begin{pmatrix} 2 & a & -1 \\ 1 & b & 3 \\ 0 & c & 2 \end{pmatrix}$. Although the entries of the second column are unknown, we are given that $|M| = 12$.

(a) Solve the system $M \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 2 \\ -2 \\ 0 \end{pmatrix}$ FOR y ONLY, using Cramer's rule.

(b) Find the value of the determinant (using the given information about (a))

$$\begin{vmatrix} 2c + a & 2 & 3 \\ 2c & 0 & 4 \\ 3a + b & 7 & 0 \end{vmatrix}$$

(c) Find the adjoint of $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$

THE UNIVERSITY OF MANITOBA

DATE: April 14, 2003

FINAL EXAMINATION

PAPER NO: 131

PAGE 2 of 8

DEPARTMENT & COURSE NO: 136.130

TIME: 2 HOURS

EXAMINATION: Vector Geometry &
Linear Algebra

EXAMINERS: Various

Values

- [20] 6. Let $\mathbf{u} = (1,1,1)$, $\mathbf{v} = (-1,-2,3)$ and $\mathbf{w} = (2,2,0)$. Find:
- $\|2\mathbf{u} - 3\mathbf{v}\|$.
 - the area of the triangle with vertices at $A(1,1,1)$, $B(-1,-2,3)$, $C(2,2,0)$.
 - the volume of the parallelepiped, determined by \mathbf{u} , \mathbf{v} and \mathbf{w} .
 - show that two of \mathbf{u} , \mathbf{v} and \mathbf{w} are orthogonal.
 - find the projection of \mathbf{w} onto \mathbf{u} .
- [8] 7. Consider the vectors \mathbf{u} , \mathbf{v} , where $\|\mathbf{u}\| = 2$, $\|\mathbf{v}\| = 3$, and the angle between \mathbf{u} and \mathbf{v} is $\theta = \frac{\pi}{3}$ radians ($= 60^\circ$). Find
- $\mathbf{u} \cdot \mathbf{v}$.
 - $\|\mathbf{u} \times \mathbf{v}\|$.
- [12] 8. Let P be the plane whose equation is $x + 3y - z = 2$. Define the points $A(5,0,3)$, $B(2,-1,3)$, $C(1,1,2)$ and $D(0,0,1)$.
- Find a vector normal to the plane.
 - Show that A and C are on the plane P but B and D are not.
 - Find parametric equations for the line orthogonal to P and passing through A .
- [10] 9. Consider the vector space $P_2 = \{a + bx + cx^2 \mid a, b, c \in \mathbf{R}\}$ and the subsets $S = \{f(x) \in P_2 \mid f(2) = 0\}$ and $T = \{a + bx + cx^2 \in P_2 \mid a, b, c \geq -1\}$.
- Use the subspace test to show that S is a subspace of P_2 .
 - Show that T is not a subspace of P_2 , by giving a specific case for which part of the subspace test fails.
- [5] 10. Define "linearly independent set".
- [12] 11. Let $A = \begin{pmatrix} 1 & 2 & 3 & 1 \\ 1 & 2 & 3 & 2 \\ 2 & 4 & 6 & 0 \\ 1 & 2 & 2 & 1 \end{pmatrix}$.
- Find a basis for, and dimension of, the row space of A .
 - Find a basis for, and dimension of, the column space of A .
 - Find a basis for, and dimension of, the solution space of the system $A\mathbf{x} = \mathbf{0}$.